

Recent Trials of Second-Line ART: Lessons Learned & Applications to Clinical Practice

Brian R. Wood, MD

**Associate Professor of Medicine, University of Washington
Medical Director, MWAETC Project ECHO**

Last Updated: August 25, 2022

Disclosures

No conflicts of interest or relationships to disclose.

Disclaimer

Funding for this presentation was made possible by U1OHA29296 from the Human Resources and Services Administration HIV/AIDS Bureau. The views expressed do not necessarily reflect the official policies of the Department of Health and Human Services nor does mention of trade names, commercial practices, or organizations imply endorsement by the U.S. Government. *Any trade/brand names for products mentioned during this presentation are for training and identification purposes only.*

Case

- 52-year-old cisgender man with longstanding HIV
- Viral load suppressed on rilpivirine/TAF/FTC for several years
- Prior ART: efavirenz/TDF/FTC
- Lapse in adherence following onset of COVID-19 pandemic
- Viral load rebound to 1,250 copies/mL
- Genotype: new E138K and M184V mutations
- *Which new ART regimen would you recommend?*

Background

- Following virologic failure of first-line ART, traditional practice was to aim for 3 fully active drugs in new regimen
- In areas without access to resistance testing, new regimen often included a boosted PI plus switch from TDF to AZT
- With widespread availability of dolutegravir, is this still the optimal strategy?
- As many countries roll out tenofovir DF-lamivudine-dolutegravir (“TLD”) as first-line ART, can it also be offered as empiric second-line ART?

DAWNING

Dolutegravir (DTG) vs. ritonavir-boosted lopinavir (LPV/r), each with two NRTIs, following virologic failure of first-line ART

DTG vs LPV/r, each with two NRTIs, after first-line ART virologic failure

DAWNING: Design

- **Design**

- Open-label, randomized, phase 3b study performed in multiple countries in sub-Saharan Africa, South America, Central America, Asia, and Eastern Europe

- **Including Criteria**

- Age ≥ 18 with HIV-1
- Virologic failure after at least 6 months taking NNRTI plus 2 NRTIs (HIV RNA ≥ 400 copies/mL x 2)
- No history of taking a boosted PI or INSTI
- All had genotype resistance test at baseline
- All received investigator-selected NRTIs (at least one fully active based on genotype)

DTG + 2 NRTIs
(n = 312)

LPV/r + 2 NRTIs
(n = 312)

DTG vs LPV/r, each with two NRTIs, after first-line ART virologic failure

DAWNING: Results

Baseline Characteristics	DTG + 2 NRTIs (n = 312)	LPV/r + 2 NRTIs (n = 312)
Age, years, mean (range)	37.5 (19-64)	38.7 (18-72)
Male sex, n (%)	196 (63)	209 (67)
CD4 T cell count, mean (SD), log ₁₀ cells/mm ³	2.1 (0.5)	2.2 (0.4)
CD4 T cell count <200 cells/mm ³	166 (53%)	151 (48%)
Mean HIV RNA, mean (SD), log ₁₀ copies/mL	4.2 (0.9)	4.2 (0.9)
HIV RNA >100,000 copies/mL, n (%)	70 (22)	63 (20)
Duration of previous ART, median (IQR), weeks	86.4 (48.4-230.9)	90.9 (45.0-199.5)
Prior NNRTI: efavirenz	242 (78%)	242 (78%)
Prior NNRTI: nevirapine	70 (22%)	69 (22%)
Prior NNRTI: rilpivirine	0	1 (<1%)

DTG vs LPV/r, each with two NRTIs, after first-line ART virologic failure

DAWNING: Results

Baseline Characteristics	DTG + 2 NRTIs (n = 312)	LPV/r + 2 NRTIs (n = 312)
NRTIs in new regimen		
TDF/FTC or TDF/3TC	132 (42%)	121 (39%)
AZT/3TC	128 (41%)	134 (43%)
TDF + AZT	36 (12%)	41 (13%)
Fully susceptible NRTIs in new regimen		
0 to <1	30 (10%)	36 (12%)
1 to 2	221 (71%)	212 (68%)
2	61 (20%)	64 (21%)

DTG vs LPV/r, each with two NRTIs, after first-line ART virologic failure


DAWNING: Results

Baseline Characteristics (NRTI Resistance Mutations)	DTG + 2 NRTIs (n = 312)	LPV/r + 2 NRTIs (n = 312)
K65R	95 (30%)	92 (29%)
K70E	33 (11%)	37 (12%)
M184V/I only	77 (25%)	85 (27%)
M184V/I plus other major NRTI mutation	184 (59%)	167 (54%)
Other major NRTI mutation	90 (29%)	88 (28%)
Thymidine analog mutation (TAMs)	71 (23%)	81 (26%)

DTG vs LPV/r, each with two NRTIs, after first-line ART virologic failure

DAWNING: Results


Virologic response at 48 weeks (intention-to-treat analysis), stratified by baseline viral load (VL)



DTG vs LPV/r, each with two NRTIs, after first-line ART virologic failure

DAWNING: Results


Virologic response at 48 weeks, stratified by M184V and use of XTC (FTC or 3TC)



DTG vs LPV/r, each with two NRTIs, after first-line ART virologic failure

DAWNING: Results

Virologic response at 48 weeks, stratified by number of fully active NRTIs



DTG vs LPV/r, each with two NRTIs, after first-line ART virologic failure

DAWNING: Results

Cases of virologic failure with emergent dolutegravir resistance

DTG arm: 11 participants met criteria for virologic failure (VF); 2 had emergent DTG resistance
LPV/r arm: 30 met criteria for VF; 3 emergent NRTI resistance, zero emergent PI resistance

HIV Subtype	Study NRTIs	Baseline HIV RNA (copies/mL)	HIV RNA at Virologic Failure (copies/mL)	Baseline NRTI RAM(s)	Emergent INSTI RAM(s)	Emergent NRTI RAM(s)
B	TDF/FTC	461,000	2,464	M184V + K219K/E	G118R	D67N
C	AZT/3TC	1.2 million	454	M184V + K70E	Multiple	None

DAWNING: Conclusions and Limitations

- After 48 weeks, DTG plus 2 NRTIs showed superior efficacy compared to LPV/r plus 2 NRTIs following virologic failure on NNRTI plus 2 NRTIs
- Supports DTG + 2 NRTIs as second-line ART, even if only one NRTI fully active
- Limitations: open-label, baseline resistance testing performed (not generalizable to low-resource settings), use of LPV/r as comparator, use of TDF (not TAF)
- Outstanding questions: efficacy of TDF/FTC versus AZT/3TC in new regimen, efficacy and safety of switching in the absence of genotype

NADIA

**Dolutegravir (DTG) vs. ritonavir-boosted darunavir (DRV/r),
each with TDF/3TC or AZT/3TC, following virologic failure on
first-line ART**

DTG vs DRV/r, each with TDF/3TC or AZT/3TC, after first-line ART virologic failure

NADIA: Design

- **Design**

- Open-label, prospective, multicenter, two-by-two factorial, randomized, non-inferiority, 96-week trial conducted in Uganda, Kenya, and Zimbabwe

- **Including Criteria**

- Age ≥ 12 with HIV-1
- Virologic failure after at least 6 months taking NNRTI plus 2 NRTIs (HIV RNA $\geq 1,000$ copies/mL twice)
- No history of taking a boosted PI or INSTI
- Nurse-led visits, standard of care visit frequency, emphasis on adherence counseling
- No genotype at enrollment

DTG + TDF/3TC

(n = 118)

DTG + AZT/3TC

(n = 117)

DRV/r + TDF/3TC

(n = 115)

DRV/r + AZT/3TC


(n = 114)

DTG vs DRV/r, each with TDF/3TC or AZT/3TC, after first-line ART virologic failure NADIA: Results

Baseline Characteristics	DTG (n = 235)	DRV/r (n = 229)
Age, years, median (IQR)	33 (28-40)	35 (28-42)
Female sex, n (%)	140 (59.6)	142 (62.0)
CD4 T cell count, median (IQR), cells/mm ³	189 (58-388)	202 (84-357)
CD4 T cell count <200 cells/mm ³	125 (53.2)	113 (49.3)
Median HIV RNA (IQR), log ₁₀ copies/mL	4.5 (3.9-5.1)	4.4 (3.8-5.1)
HIV RNA ≥100,000 copies/mL, n (%)	66 (28.1)	62 (27.1)
Duration of previous ART, median (IQR), years	3.6 (1.4-6.3)	3.7 (1.7-5.9)
K65R present at baseline	120 (52.6)	106 (47.1)
M184V/I present at baseline	196 (86.0)	195 (86.7)


DTG vs DRV/r, each with TDF/3TC or AZT/3TC, after first-line ART virologic failure NADIA: Results

Virologic response at 48 weeks (by intention-to-treat analysis)




DTG vs DRV/r, each with TDF/3TC or AZT/3TC, after first-line ART virologic failure NADIA: Results

Virologic response at 48 weeks, stratified by number of predicted active NRTIs



DTG vs DRV/r, each with TDF/3TC or AZT/3TC, after first-line ART virologic failure NADIA: Results

Virologic response at week 48, stratified by NRTI backbone (TDF/3TC or AZT/3TC)



NADIA

Week 96 results presented at CROI 2022

- DTG + 2 NRTIs remained non-inferior to DRV/RTV + 2 NRTIs
- TDF/3TC now superior efficacy compared to AZT/3TC
- 9 cases of emergent DTG resistance (6 in AZT/3TC group, 3 TDF/FTC)
- Conclusions:
 - DTG + 2 NRTIs can be used as second-line ART, even if NRTIs predicted to have limited or no activity, but emergent DTG resistance may be a concern
 - DRV/r + 2 NRTIs has efficacy equivalent to DTG + 2 NRTIs in second-line treatment, without concern for resistance

- How should we apply these results to clinical practice?
 - DTG (or BIC) or DRV/r effective with <2 fully active NRTIs (e.g. with M184V); is this the end of AZT once and for all as part of HIV treatment?
 - Would you offer DTG (or BIC) plus TDF/FTC or TAF/FTC in the setting of M184V and K65R? Or M184V + TAMs? Are you comfortable if there is <1 active NRTI?
 - Cases of intermediate-to-high-level resistance to DTG occurred in NADIA; should this give us pause? Should we opt for DRV/r instead of DTG in certain scenarios?
- Returning to case, with M184V and viral load 1,250 copies/mL, which regimen would you recommend? What if the viral load were 12k copies? Or 100k copies?

Acknowledgment

This Mountain West AIDS Education and Training (MWAETC) program is supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) as part of an award totaling \$3,098,654 with 0% financed with non-governmental sources.

The content in this presentation are those of the author(s) and do not necessarily represent the official views of, nor an endorsement by, HRSA, HHS, or the U.S. Government.

