Texas Billing and Coding Guide

A Resource for Providers of HIV, STD and Related Services





Texas Billing and Coding Guide

A Resource for Providers of HIV, STD and Related Services

ACKNOWLEDGEMENTS

Cardea

For more than 40 years, Cardea has provided training, organizational capacity building, and research and evaluation services to improve organizations' abilities to deliver accessible, culturally proficient, and compassionate services to their clients. Cardea serves as the STD-related Reproductive Health Training and Technical Assistance Center (STDRHTTAC) for U.S. Public Health Regions VI, IX, and X. Cardea has developed this guide in partnership with the Texas Oklahoma AIDS Education and Training Center as part of a resource portfolio to support public health programs with third-party billing for sexually transmitted disease (STD) and other related services. Along with this and other resources, the portfolio will include:

- Webinars and other resource materials
- An online learning community to facilitate peer learning
- Customized training and technical assistance

Contact us for more information:

Nikki Trevino, ntrevino@cardeaservices.org or 512-474-2166

Cardeaservices.org

Funded by a cooperative agreement by the Office of Population Affairs, within the Office of the Assistant Secretary for Health in collaboration with the Division of STD Prevention within the Centers for Disease Control and Prevention, National Center for HIV/AIDS, Viral Hepatitis, STD, and TB Prevention.

Texas/ Oklahoma AIDS Education and Training Center

The Texas/Oklahoma AIDS Education and Training Centers is one of the regional offices of the AIDS Education and Training Center (AETC) Program. The AETC Program is the professional training arm of the Ryan White HIV/AIDS Program. One of the largest and most comprehensive professional education programs dedicated to HIV/AIDS, the AETC Program was developed by Health Resources and Services Administration (HRSA), U.S. Public Health Service, and the Department of Health and Human Services to provide customized, multi-disciplinary training programs for healthcare providers treating persons living with HIV/AIDS.

Contact us for more information:

Marcos Alcorn, marcos.alcorn@phhs.org or 214-590 - 1650

Aidseducation.org

This Texas Billing and Coding Guide was made possible by AETC grant award #H4AHA00061 from the HIV/AIDS Bureau of the Health Resources Services Administration (HRSA), U.S. Department of Health and Human Services (HHS). Dallas County Hospital District operates an AIDS Education and Training Center (AETC) that strengthens the capacity of health care professionals to care for people living with HIV/AIDS through training and technical assistance. The information presented here is the consensus of HIV/AIDS specialists in the Texas/Oklahoma AIDS Education & Training Center (TX/OK AETC) and does not necessarily represent the official views of HRSA/HAB.

DISCLAIMER

This guide was prepared as a service to the public and is only intended to be a general summary. It is not intended to take the place of either the written law or regulations. We encourage readers to review the specific statutes, regulations and other interpretive materials for a full and accurate statement of their contents. Specific coding and payer guidelines should be reviewed prior to the submission of claims for reimbursement.

OVERVIEW

This guide is designed to be a resource for programs providing HIV, STD and related services and to assist in the transition from ICD-9 to ICD-10 coding. It includes basic information about HIV and STD screening and testing, an overview of coding guidelines, and common ICD-9, ICD-10, and CPT codes. Several coding scenarios are included along with accompanying resources and references. While this is not an exhaustive list of codes for HIV, STD and related services and diagnoses, these common codes offer a quick reference to help improve coding efficiency. Always follow current legal guidelines and standards of practice when providing medical services.

Coding for HIV, STD and related services is an essential practice for programs that are preparing for billing third party payers. Beginning to properly code for services is a critical step in improving revenue cycle management and developing sustainable systems. For more information about billing and revenue cycle management, visit the resources section of this document.

SCREENING RECOMMENDATIONS

Third-party payer coverage for CDC recommended screening services may vary. Contact payers to determine billing eligibility.

HIV Recommendations

The objectives of the following recommendations are to: increase HIV screening of patients, including pregnant women, in health care settings; foster earlier detection of HIV infection; identify and counsel persons with unrecognized HIV infection and link them to clinical and prevention services; and further reduce perinatal transmission of HIV in the United States.

The Centers for Disease Control and Prevention (CDC) recommendations for patients in all health care settings:

- HIV screening is recommended for patients in all health care settings after the patient is notified that testing will be performed unless the patient declines (opt-out screening).
- Persons at high-risk for HIV infection should be screened for HIV at least annually.
- Separate written consent for HIV testing should not be required. General consent for medical care should be considered sufficient to encompass consent for HIV testing.
- Prevention counseling should not be required with HIV diagnostic testing or as part of HIV screening programs in health care settings.

Why provide routine screening for HIV/AIDS?

- Twenty percent (20%) of people living in the U.S. with HIV do not know they are infected.
- HIV is a serious health condition that can be reliably diagnosed prior to symptoms development, and patients are more likely to be tested if it is suggested by a clinician.

- Routine screening helps to de-stigmatize HIV.
- Early entry into care increases the likelihood of a longer, healthier life. Unfortunately, an estimated 39% of people with HIV in the U.S are not diagnosed until they are in the later stages of the disease.
- Transmission rates are higher in people who do not know they have HIV. Individuals who know that they are HIV positive are more likely to engage in risk reduction efforts.
- Appropriate treatment during pregnancy can reduce the perinatal transmission rate to < 2%.
 Pregnant women who know they are infected are better able to make critical decisions about care for themselves and their infants.

Source: Centers for Disease Control and Prevention

STD Recommendations

The Centers for Disease Control and Prevention makes the following recommendations for STD screening:

- Chlamydia—screen women under age 25 and others, including men, at increased risk
- Gonorrhea—screen women at increased risk
- Syphilis—screen women exposed to syphilis
- Hepatitis B—provide prevaccination screening for women at increased risk
- Hepatitis C— screen women at risk
- Herpes Simplex Virus—do not screen general population

Other health organizations also provide screening recommendations for STD that may vary slightly. For a list comparing these recommendations, please see the resources section.

Why provide screening for STDs?

- Chlamydia and Gonorrhea prevalence is highest among adolescents and young adults under 25
- STDs can have severe effects to pregnant women, their partners and fetuses
- STDs can increase the risk of contracting HIV

HIV/STD Recommendations for Pregnant Women

For pregnant women, the CDC recommends the following screening:

- STD and HIV screening should be included in the routine panel of prenatal screening tests for all pregnant women in the first trimester or prenatal appointment.
- STD screening for pregnant women should include the following: syphilis, hepatitis B, hepatitis C (increased risk patients only), chlamydia, and gonorrhea (increased risk patients only).
- STD and HIV screening is recommended after the patient is notified that testing will be performed unless the patient declines (opt-out screening).
- Separate written consent for HIV testing should not be required; general consent for medical care should be considered sufficient to encompass consent for HIV testing.
- Repeat screening in the third trimester is recommended in certain jurisdictions with elevated rates
 of HIV infection among pregnant women. Women at increased risk for HIV and STDs should also be
 screened for syphilis, HIV, chlamydia and gonorrhea in the third trimester.

Source: Centers for Disease Control and Prevention

CODING GUIDELINES

What is documentation and why is it important?

According to the Centers for Medicare and Medicaid Services (CMS), medical record documentation is required to record pertinent facts, findings, and observations about an individual's health history including past and present illnesses, examinations, tests, treatments, and outcomes. The medical record chronologically documents the care of the patient and is an important element contributing to high quality care. The medical record facilitates:

- The ability of the physician and other health care professionals to evaluate and plan the patient's immediate treatment, and to better manage the patient's care over time
- Communication and continuity of care among physicians and other health care professionals involved in the patient's care
- Accurate and timely claims review and payment
- Appropriate utilization review and quality of care evaluations and
- Collection of data that may be useful for research and education

Document every step you take. Remember, if it's not documented in the record, it did not happen.

Source: Centers for Medicare and Medicaid Services

International Classification of Diseases Diagnosis Codes

The International Classification of Diseases (ICD) is a system of coding maintained by the World Health Organization that is used to describe diseases, symptoms, abnormal findings, and external causes of injury.

On October 1, 2014, the ICD-9 code sets used to report medical diagnoses and inpatient procedures will be replaced by ICD-10 code sets. The transition to ICD-10 is required for everyone covered by the Health Insurance Portability and Accountability Act (HIPAA). Please note that the change to ICD-10 does not affect CPT coding for outpatient procedures and physician services. Included below are common ICD codes that are used for coding STD and HIV services as reference. Multiple codes may need to be documented to support medical necessity when billing for services. ICD coding guidelines, including specific rules for coding HIV, should always be consulted for accurate coding purposes.

ICD-9 to ICD-10 Crosswalk

| | HIV Specific Codes | | | |
|--------|---|--------|---|--|
| ICD-9 | Description | ICD-10 | Description | |
| V70.0 | Routine general medical examination at a health care facility | Z00.00 | Encounter for general adult medical examination without abnormal findings | |
| | | Z00.01 | Encounter for general adult medical examination abnormal findings* | |
| | | | *Use additional code to identify abnormal findings (R70-R94) | |
| V73.89 | Special screening examination for other specified viral diseases (HIV/AIDS) | Z11.4 | Encounter for screening for human immunodeficiency virus (HIV) | |
| V65.44 | HIV counseling | Z71.7 | HIV counseling | |
| V69.2 | High-risk sexual behavior | Z72.51 | High-risk heterosexual behavior | |
| | | Z72.52 | High-risk homosexual behavior | |
| | | Z72.53 | High-risk bisexual behavior | |
| V69.8 | Other problems related to lifestyle | Z72.89 | Other problems related to lifestyle (self-damaging behavior) | |
| 042 | HIV disease | B20 | HIV disease *Code first HIV disease complicating pregnancy, childbirth and the puerperium, if applicable (098.7-) *Use additional code(s) to identify all manifestations of HIV infection | |

| ICD- 9 | Description | ICD- 10 | Description |
|--------|--|---------------|--|
| 079.53 | HIV, type 2 (HIV-2) * Report as secondary diagnosis code only (when applicable) | B97.35 | HIV, type 2 (HIV-2) as the cause of diseases classified elsewhere |
| V08 | Asymptomatic HIV infection status | Z21 | Asymptomatic HIV infection status. |
| | | | * Code first HIV disease complicating pregnancy, childbirth and the puerperium, if applicable (098.7-) |
| | Screening Tests: | Increased | Risk of Infection |
| V70.0 | Routine general medical examination at a health care facility | Z00.00 | Encounter for general adult medical examination without abnormal findings |
| | | Z00.01 | Encounter for general adult medical examination abnormal findings |
| | | | * Use additional code to identify abnormal findings (R70-R94) |
| V01.1 | Contact with or exposure to tuberculosis | Z20.1 | Contact with and (suspected) exposure to tuberculosis |
| V01.6 | Contact with or exposure to venereal diseases | Z20.2 | Contact with and (suspected) exposure to infections with a predominantly sexual mode of transmission |
| V02.7 | Carrier or suspected carrier of gonorrhea | Z22.4 | Carrier of infections with a predominantly sexual mode of transmission |
| V02.8 | Carrier or suspected carrier of other venereal diseases | | |
| V12.00 | Personal history of unspecified infectious and parasitic disease | Z86.19 | Personal history of other infectious and parasitic diseases |
| V12.09 | Personal history of other infectious and parasitic diseases | | *Conditions classifiable to A00-B89, B99, |
| V69.2 | High-risk sexual behavior | Z72.51 | High-risk heterosexual behavior |
| | | Z72.52 | High-risk homosexual behavior |
| | | Z72.53 | High-risk bisexual behavior |
| V69.8 | Other problems related to lifestyle | Z72.89 | Other problems related to lifestyle |
| | | | (Self-damaging behavior) |
| V73.81 | Special screening examination for human papillomavirus (HPV) | Z11.51 | Encounter for screening for HPV |
| V73.88 | Special screening examination for other specified chlamydial diseases | Z11.8 | Encounter for screening for other infectious and parasitic diseases |
| V73.98 | Special screening examination for unspecified chlamydial disease | | (Encounter for screening for chlamydia, rickettsial, spirochetal or mycoses) |

| V73.89 | Special screening examination for other specified viral diseases (HIV) | Z11.59 Z11.4 | Encounter for screening for other viral diseases Encounter for screening for HIV | |
|--------|--|-----------------|---|--|
| V73.99 | Special screening examination for unspecified viral disease | | Encounter for screening for other viral diseases | |
| V74.1 | Screening examination for pulmonary tuberculosis | Z11.1 | Encounter for screening for respiratory tuberculosis | |
| V74.5 | Screening examination for venereal disease | Z11.3 | Encounter for screening for infections with a predominantly sexual mode of transmission | |
| V75.8 | Screening examination for other specified parasitic infections | Z11.0 | Encounter for screening for intestinal infectious diseases | |
| | | Z11.8 | Encounter for screening for other infectious and parasitic diseases | |

Diagnostic Testing

| ICD- 9 | Description | ICD- 10 | Description | |
|------------------|--|---------|---|--|
| 053.9 | Herpes zoster without mention of complication | B02.9 | Zoster without complications (Shingles, zona) | |
| 054.10 | Genital herpes, unspecified | A60.9 | Anogenital herpesviral infection, unspecified | |
| 054.11 054.12 | Herpetic vulvovaginitis Herpetic ulceration of vulva A60.04 | | Herpesviral vulvovaginitis (Herpesviral (herpes simplex) ulceration, vaginitis, vulvitis) | |
| 054.13 | Herpetic infection of penis A60.01 | | Herpesviral infection of penis | |
| 054.19 | Other genital herpes | A60.09 | Herpesviral infection of other urogenital tract | |
| 054.2 | Herpetic gingivostomatitis (Simplex Oral) | B00.2 | Herpesviral gingivostomatitis and pharyngotonsillitis | |
| 054.8 054.9 | Herpes simplex with unspecified complication Herpes simplex without mention of complication | | Herpesviral infection, unspecified (Herpes simplex infection NOS) | |
| 078.0 | Molluscum contagiosum | B08.1 | Molluscum contagiosum | |
| 078.11 | Condyloma acuminatum | A63.0 | Anogenital (venereal) warts (Anogenital warts due to HPV, Condyloma acuminatum) | |

| ICD- 9 | Description | ICD- 10 | Description |
|--------|---|----------|--|
| 078.19 | Other specified viral warts | B07.8 | Other viral warts |
| 078.88 | Other specified diseases due to chlamydiae | A74.89 | Other chlamydial diseases |
| 079.98 | Unspecified chlamydial infection | A74.9 | Chlamydial infection, unspecified |
| | | | (Chlamydiosis NOS) |
| 091.0 | Genital syphilis (primary) | A51.0 | Primary genital syphilis |
| | | | (Syphilitic chancre NOS) |
| 091.1 | Primary anal syphilis | A51.1 | Primary anal syphilis |
| 091.4 | Adenopathy due to secondary syphilis | A51.49 | Other secondary syphilitic conditions |
| 091.69 | Secondary syphilis of other viscera | | (Secondary syphilitic lymphadenopathy or |
| 091.7 | Secondary syphilis, relapse | | myositis) |
| 091.89 | Other forms of secondary syphilis | | |
| 091.9 | Unspecified secondary syphilis | | |
| 092.0 | Early syphilis, latent, serological relapse after treatment. | A51.5 | Early syphilis, latent |
| 092.9 | Early syphilis, latent, unspecified | | (Syphilis (acquired) without clinical manifestations, with positive serological reaction and negative spinal fluid test, less |
| | | | than 2 years after infection.) |
| 096 | Late syphilis, latent | A52.8 | Late syphilis, latent |
| | | | (Syphilis (acquired) without clinical manifestations, with positive serological reaction and negative spinal fluid test, 2 years or more after infection.) |
| 097.9 | Syphilis, unspecified | A53.9 | Syphilis, unspecified |
| | | | (Infection due to Treponema pallidum NOS, Syphilis (acquired) NOS) |
| 098.0 | Gonococcal infection (acute) of lower genitourinary tract | A54.00 | Gonococcal infection of lower genitourinary tract, unspecified |
| 098.2 | Gonococcal infection, chronic, of lower genitourinary tract | | |
| 098.10 | Gonococcal infection (acute) of upper genitourinary tract, site unspecified | A54.29 | Other gonococcal genitourinary infections |
| 098.19 | Gonococcal (acute) upper genitourinary NEC | | |
| 098.12 | Gonococcal prostatitis acute | A54.22 | Gonococcal prostatitis |
| 098.32 | Gonococcal prostatitis, chronic | | |
| | | <u> </u> | |

| ICD-9 | Description | ICD - 10 | Description |
|--------|---|------------------------------|---|
| 098.13 | Gonococcal epididymo-orchitis (acute) | A54.23 | Gonococcal infection of other male genital |
| 098.14 | Gonococcal seminal vesiculitis (acute) | | organs |
| 098.33 | Gonococcal orchitis (chronic) | | (Gonococcal epididymitis, Gonococcal orchitis) |
| 098.34 | Gonococcal seminal vesiculitis (chronic) | | |
| 098.15 | Gonococcal cervicitis (acute) | A54.03 | Gonococcal cervicitis, unspecified |
| 098.35 | Gonococcal cervicitis (chronic) | | |
| 098.16 | Gonococcal endometritis (acute) | A54.24 | Gonococcal female pelvic inflammatory disease |
| | | | (Gonococcal pelviperitonitis) |
| 098.7 | Gonococcal infection of anus and rectum | A54.6 | Gonococcal infection of anus and rectum |
| 099.0 | Chancroid | A57 | Chancroid (Ulcus molle) |
| 099.1 | Lymphogranuloma venereum | A55 | Chlamydial lymphogranuloma (venereum) |
| 099.2 | Granuloma inguinale | A58 | Granuloma inguinale |
| 099.40 | Other nongonococcal urethritis, | N34.1 | Nonspecific urethritis |
| 099.41 | unspecified Other nongonococcal urethritis, chlamydia trachomatis | | (Nongonococcal urethritis, Nonvenereal urethritis) |
| 099.52 | Other venereal diseases due to chlamydia trachomatis, anus and rectum | A56.3 | Chlamydial infection of anus and rectum |
| 099.53 | Other venereal diseases due to chlamydia trachomatis, lower genitourinary sites | A56.00 | Other venereal diseases due to chlamydia trachomatis, lower genitourinary sites |
| 110.3 | Dermatophytosis of groin and perianal | B35.6 | Tinea cruris |
| | area | | (DHOBI Itch, Groin ringworm, jock itch) |
| 112.0 | Candidiasis of mouth (Thrush Oral) | B37.0 | Candidal stomatitis |
| | | B37.83 | Candidal cheilitis |
| 112.1 | Candidiasis of vulva and vagina | B37.3 | Candidiasis of vulva and vagina |
| | (Candidal vulvovaginitis, moni vulvovaginitis, vaginal thrush) | | (Candidal vulvovaginitis, monilial vulvovaginitis, vaginal thrush) |
| 112.84 | Candidal esophagitis | B37.81 | Candidal esophagitis |
| 131.01 | Trichomonal vulvovaginitis | A59.01 | Trichomonal vulvovaginitis |
| 133.0 | Scabies | B86 Scabies (Sarcoptic itch) | |

| ICD- 9 | Description | ICD- 10 | Description |
|----------------|---|---------|--|
| 599.0 | Urinary tract infection (UTI), site not | N39.0 | Urinary tract infection, site not specified |
| | specified | | *Use Additional code (B95-B97), to identify infectious agent |
| 614.3 | Acute parametritis and pelvic cellulitis | N73.0 | Acute parametritis and pelvic cellulitis |
| | | | (Abscess of broad ligament or parametrium, Pelvic cellulitis female) |
| | | | *Use Additional code (B95-B97), to identify infectious agent |
| 616.0 | Cervicitis and endocervicitis | N72 | Inflammatory disease of cervix uteri |
| | | | (Cervicitis (with or without) erosion or ectropion; Endocervicitis (with or without) erosion or ectropion; Exocervicitis (with or without) erosion or ectropion) |
| | | | *Use Additional code (B95-B97), to identify infectious agent |
| 616.10 | Vaginitis and vulvovaginitis, | N76.0 | Acute vaginitis |
| | unspecified | N76.1 | Subacute and chronic vaginitis |
| | | N76.2 | Acute vulvitis |
| | | N76.3 | Subacute and chronic vulvitis |
| | | | *Use Additional code (B95-B97), to identify infectious agent |
| 623.5 623.6 | Leukorrhea, not specified as infective Vaginal hematoma | N89.8 | Other specified non-inflammatory disorders of vagina |
| 623.8 | Other specified non-inflammatory disorders of vagina | | (Leukorrhea NOS, Old vaginal laceration, Pessary ulcer of vagina) |
| 782.1 | Rash and other nonspecific skin eruption | R21 | Rash and other nonspecific skin eruption |
| 788.1 | Dysuria | R30.0 | Dysuria |
| | | R30.9 | Painful micturition, unspecified |
| | | | (Painful urination NOS) |
| 788.7 | Urethral discharge | R36.0 | Urethral discharge without blood |
| | | R36.9 | Urethral discharge, unspecified |
| | | | (Penile discharge NOS, Urethrorrhea) |
| 795.00 | Abnormal glandular Papanicolaou smear of cervix | R87.619 | Unspecified abnormal cytological findings in specimens from cervix uteri |

| ICD- 9 | Description | ICD- 10 | Description |
|------------------|--|---------|--|
| 795.10 795.19 | Abnormal glandular Papanicolaou smear of vagina Other abnormal Papanicolaou smear of vagina and vaginal HPV | R87.628 | Other abnormal cytological findings on specimens from vagina * Use additional code to identify acquired absence of the uterus and cervix, if applicable |
| | vagilia aliu vagiliai i ir v | | (Z90.71-) |
| V05.3 | Need for prophylactic vaccination and inoculation against viral hepatitis | Z23 | Encounter for immunization |
| V05.8 | Need for prophylactic vaccination and inoculation against other specified disease | | |
| V25.03 | Encounter for emergency contraceptive counseling and prescription | Z30.012 | Encounter for prescription of emergency contraception |
| V25.09 | Other general counseling and advice on contraceptive management | Z30.09 | Encounter for other general counseling and advice on contraception |
| V65.42 | Counseling on substance use and abuse | Z71.41 | Alcohol abuse counseling and surveillance of alcoholic |
| V72.41 | Pregnancy exam or test, negative result | Z32.02 | Encounter for pregnancy test, result negative |
| V72.42 | Pregnancy exam or test, positive result | Z32.01 | Encounter for pregnancy test, result positive |

Pregnancy Related Codes

| Supervision of normal first pregnancy | Z34.00 | Encounter for supervision of normal first pregnancy unspecified trimester |
|---------------------------------------|--------------------------------------|---|
| | Z34.01 Z34.02 Z34.03 | first trimester second trimester third trimester |
| Supervision of other normal pregnancy | Z34.80 Z34.81 Z34.82 Z34.83 | Encounter for supervision of other normal pregnancy unspecified trimester first trimester second trimester third trimester |
| Other high-risk pregnancy | 009 | Supervision of high-risk pregnancy (requires 4th and 5th digits) |
| | 009.0 009.00 009.01 009.02 | Supervision of pregnancy with history of infertility unspecified trimester first trimester second trimester third trimester |
| | | Z34.02 Z34.03 upervision of other normal pregnancy Z34.8 Z34.80 Z34.81 Z34.82 Z34.83 Uther high-risk pregnancy 009 009.0 009.0 009.00 009.01 |

| ICD- 9 | Description | ICD- 10 | Description |
|--------|---------------------------------------|--|---|
| V23.8 | Other high-risk pregnancy (continued) | 009.10 009.11 009.12 009.13 | Supervision of pregnancy with history of ectopic or molar pregnancy unspecified trimester first trimester second trimester third trimester |
| | | 009.2 | Supervision of pregnancy with other poor reproductive or obstetric history |
| | | 009.21 009.211 009.212 009.213 009.219 | Supervision of pregnancy with history of preterm labor first trimester second trimester third trimester unspecified trimester |
| | | 009.29 009.291 009.292 009.293 009.299 | Supervision of pregnancy with other poor reproductive or obstetric history first trimester second trimester third trimester unspecified trimester |
| | | 009.3 009.30 009.31 009.32 009.33 | Supervision of pregnancy with insufficient antenatal care unspecified trimester first trimester second trimester third trimester |
| | | 009.4 009.40 009.41 009.42 009.43 | Supervision of pregnancy with grand multiparity unspecified trimester first trimester second trimester third trimester |
| | | 009.5 | Supervision of elderly primigravida and multigravida |
| | | 009.51 009.511 009.512 009.513 009.519 | Supervision of elderly primigravida first trimester second trimester third trimester unspecified trimester |
| | | 009.52 009.521 009.522 009.523 009.529 | Supervision of elderly multigravida first trimester second trimester third trimester unspecified trimester |

| ICD- 9 | Description | ICD- 10 | Description |
|--------|---------------------------------------|---|--|
| V23.8 | Other high-risk pregnancy (continued) | 009.6 | Supervision of young primigravida and multigravida |
| | | 009.61 009.611 009.612 009.613 009.619 | Supervision of young primigravida first trimester second trimester third trimester unspecified trimester |
| | | 009.62 09.621 009.622 009.623 009.629 | Supervision of young multigravida first trimester second trimester third trimester unspecified trimester |
| | | 009.7 009.70 009.71 009.72 009.73 | Supervision of high-risk pregnancy due to social problems unspecified trimester first trimester second trimester third trimester |
| | | 009.8 009.81 009.811 009.812 009.813 009.819 | Supervision of other high-risk pregnancies Supervision of pregnancy resulting from assisted reproductive technology first trimester second trimester third trimester unspecified trimester |
| | | 009.821 009.821 009.822 009.823 009.829 | Supervision of pregnancy with history of in utero procedure during previous pregnancy first trimester second trimester third trimester unspecified trimester |
| | | 009.89 009.891 009.892 009.893 009.899 | Supervision of other high-risk pregnancies first trimester second trimester third trimester unspecified trimester |
| | | 009.9 009.90 009.91 009.92 009.93 | Supervision of high-risk pregnancy, unspecified unspecified trimester first trimester second trimester third trimester |

Current Procedural Terminology (CPT) Codes

Current Procedural Terminology (CPT®) codes were developed and are maintained by the American Medical Association (AMA). They are a listing of standardized descriptions and five-character, alphanumeric codes that medical coders and billers use to report health care services and procedures to payers for reimbursement. The purpose of CPT® is to provide a uniform language accurately describing medical, surgical and diagnostic services. It serves as an effective means for reliable nationwide communication within the health care industry. CPT codes and guidelines should be reviewed prior to billing of services using the official AMA CPT guide. The following are a subset of common CPT codes that describe STD and HIV related diagnostic services.

Common CPT Codes

| | HIV Related Diagnostic Lab Tests | | | |
|-------|----------------------------------|---|--|--|
| CPT | Туре | Description | | |
| 86689 | Lab - HIV | Antibody; HTLV or HIV antibody, confirmatory test (eg, Western Blot) | | |
| 86701 | Lab - HIV | Antibody; HIV-1 | | |
| 86703 | Lab - HIV | Antibody; HIV-1 and HIV-2, single result | | |
| 87534 | Lab - HIV | Infectious agent detection by nucleic acid (DNA or RNA); HIV-1, direct probe technique | | |
| 87535 | Lab - HIV | Infectious agent detection by nucleic acid (DNA or RNA); HIV-1, reverse transcription and amplified probe technique | | |
| 87536 | Lab - HIV | Infectious agent detection by nucleic acid (DNA or RNA); HIV-1, reverse transcription and quantification | | |
| 87390 | Lab - HIV | Infectious agent antigen detection by enzyme immunoassay technique, qualitative or semi quantitative, multiple-step method; HIV-1 | | |
| | Oth | er STD Related Diagnostic Lab Tests | | |
| 81002 | Lab | Dipstick or tablet reagent urinalysis (non-automated) | | |
| 81003 | Lab | Dipstick or tablet reagent urinalysis (automated, without microscopy) | | |
| 81025 | Lab | Urine pregnancy test | | |
| 86317 | Lab | Hepatitis B Surface AB | | |
| 86580 | Lab | Purified protein derivative (PPD) skin test | | |
| 86592 | Lab | Syphilis Tests; Qualitative (e.g., VDRL, RPR) | | |
| 86695 | Lab | AB, Herpes Simplex Type 1 | | |

| СРТ | Туре | Description | | |
|---|-----------|---|--|--|
| 86696 | Lab | AB, Herpes Simplex Type 2 | | |
| 86803 | Lab | Hepatitis C antibody | | |
| 87077 | Lab | Neisseria Gonorrhea Culture | | |
| 87081 | Lab | Culture, presumptive, pathogenic organisms, screening only | | |
| 87205 | Lab | Smear Primary Source, Gram | | |
| 87210 | Lab | Smear, wet mount, (eg, saline, India ink, KOH preps) | | |
| 87220 | Lab | Tissue examination by KOH slide of samples from skin, hair, or nails for fungi | | |
| 87340 | Lab | Hepatitis B surface antigen | | |
| 87491 | Lab | Chlamydia trachomatis, amplified probe technique | | |
| 87591 | Lab | Neisseria Gonorrhea, amplified probe technique | | |
| 87621 | Lab | HPV, Amplified Probe Technique | | |
| 88141 | Lab | Cyto-Cytology Smear (PAP) | | |
| Vaccines - Common | | | | |
| 90746 | Toxoid | Hepatitis B Adult | | |
| 90649; 90650 | Toxoid | HPV – Gardasil; Cervarix | | |
| 90658 | Toxoid | Flu Vaccine | | |
| Administrations | | | | |
| 36415 | Admin | Collection of venous blood by venipuncture | | |
| 36416 | Admin | Collection of capillary blood specimen (eg, finger, heel, ear stick) | | |
| 90465-90474 | Admin | Administration of vaccines | | |
| 96372 | Admin | Therapeutic, prophylactic, or diagnostic injection; subcutaneous or intramuscular | | |
| | | Drugs - Common | | |
| J0561 | Drug | Injection, penicillin g benzathine, 100000 unit | | |
| J0696 | Drug | Injection, ceftriaxone sodium, per 250 mg | | |
| Procedures, Medical Visits and Other Services | | | | |
| 17110 | Procedure | Molluscum Destruction 1-14 | | |
| 17111 | Procedure | Molluscum Destruction 15+ | | |
| 46900/46924 | Procedure | Destruction of lesions, anus (simple/extensive) | | |
| 54050/54065 | Procedure | Destruction of lesions, penis (simple/extensive) | | |

| CPT | Туре | Description |
|-------------------------------|-----------|---|
| 56501/56515 | Procedure | Destruction of lesions, vulva (simple/extensive) |
| 57061 | Procedure | Destruction of lesions, vaginal (simple) |
| 55250 | Procedure | Vasectomy, unilateral or bilateral (separate procedure), including postoperative semen examination(s) |
| 99384; 99385; 99386, 99387 | E/M | Initial comprehensive preventive medicine evaluation and management, new patient; 12-17 years of age; 18-39 years of age; 40-64 years of age; 65 years and older |
| 99394; 99395; 99396; 99397 | E/M | Periodic comprehensive preventive medicine reevaluation and management, established patient; 12-17 years of age; 18-39 years of age; 40-64 years of age; 65 years and older |
| 99201 - 99205 | E/M | Office or other outpatient visit for the evaluation and management of a new patient (Brief, Focused, Expanded, Detailed, Comprehensive based on 3 key components: History, Exam and Medical-decision-making) |
| 99211 - 99215 | E/M | Office or other outpatient visit for the evaluation and management of an established patient (Brief, Focused, Expanded, Detailed, Comprehensive based on 3 key components: History, Exam and Medical-decision-making) |
| 99401-99404 | Other | Preventive counseling (*Time-based codes) |
| 99406; 99407 | Other | Smoking and tobacco-use cessation counseling visit (Intermediate > 3 minutes, <= 10 minutes; Intensive > 10 minutes) |
| 99408; 99409 | Other | Alcohol and/or substance abuse structured screening and brief intervention services; (15 to 30 minutes; >30 minutes) |
| 99051 | Other | Service(s) provided in the office during regularly scheduled evening, weekend, or holiday office hours, in addition to basic service (Also see 99050 – 99060) |
| H0049; H0050 | Other | Alcohol and/or drug screening; Alcohol and/or drug service, brief intervention, per 15 minutes |
| T1013 | Other | Interpreter Services |

Healthcare Common Procedure Coding System (HCPCS) Codes for Billing Medicare

| HCPCS | Description |
|-------|--|
| G0432 | Infectious agent antigen detection by enzyme immunoassay (EIA) technique, qualitative or semiquantitative, multiple-step method, HIV-1 or HIV-2, screening (conventional test) |
| G0433 | G0433 Infectious agent antigen detection by enzyme-linked immunosorbent assay (ELISA) technique, antibody, HIV-1 or HIV-2, screening |
| G0435 | G0435 Infectious agent antigen detection by rapid antibody test of oral mucosa transudate, HIV-1 or HIV-2, screening |
| A4267 | Contraceptive supply, condom, male, each |

Modifiers

Modifiers are two-digit codes appended to CPT and/or HCPCS codes. They provide additional information about who provided the billed procedure. You can find these modifiers in the CPT and HCPCS coding manuals. Third-party payers will also specify which modifiers you can use. In some cases, adding a modifier may directly affect payment.

CODING SCENARIOS

A **23-year-old single male** presents to your clinic wanting to know his HIV status. He declined testing and an exam. Patient met with a trained HIV counselor and was given a Rapid HIV 1:2 test using a swab of the mouth for the specimen. Results are negative. Pre and post counseling time was 10 minutes of the 15-minute encounter. What codes do we need for this visit?

CPT Codes:

99401 Preventive Medicine, Individual Counseling, 86703 for HIV-1 and HIV-2 single assay

Potential ICD-9 Codes:

V73.89 Special screening for other specified viral diseases (HIV), V65.44 HIV Counseling

A **25-year-old female** returns to your clinic for her HIV results as a follow-up to an earlier exam. The clinician advises the patient she is HIV+ (asymptomatic HIV). The clinician counsels her about what it means to be HIV positive, reviews risk factors and gives her a referral to see an HIV specialist. Face-to-face counseling with the clinician is 15 of the minutes of the 20-minute encounter. What codes do we need for this visit?

CPT Codes:

99213 for the problem focused E/M for an established patient based on time

Potential ICD-9 Codes:

V08 Asymptomatic human immunodeficiency virus [HIV] infection status, V65.44 HIV Counseling

An **18-year-old female** patient returns to your clinic Wednesday evening at 7 p.m. to have genital warts on her vulva removed that were diagnosed at her previous visit on Monday. The NP notes 20 lesions in the medical record and treats extensive lesions using cryosurgery. What codes do we need for this visit?

CPT Codes:

56515 for destruction of vulvar warts, extensive, 99051 for regularly scheduled office evening hours after 6 pm

Note: Do not bill for an E/M unless the documentation clearly supports it being separate and distinct from the lesion removal procedure

Potential ICD-9 Codes:

078.11 Condyloma accuminatum

A **26-year-old male** presents to your clinic for the first time. He meets with the physician and expresses concern over his partner having a vaginal discharge and "wants to be checked". Genital exam is negative. Urine dipstick is performed. A urine sample is obtained for GC and CT testing, and blood draw is performed for syphilis, HSV, HPV and HIV serologies. He is given condoms x 12 and counseled on safe sex practices. Face-to-face counseling time with the clinician is > 50% of the 30 minute encounter. What codes do we need for this visit?

CPT Codes:

99203 for the problem-focused E/M for a new patient based on time, 81002 for urinalysis dipstick without microscopy, 36415 for venipuncture; Outside lab will bill for ordered tests.

Supplies:

Check with payer re condoms

Potential ICD-9 Codes:

V01.6 Contact with or exposure to venereal diseases, V74.5 Screening Venereal Disease, V73.89 Special screening for other specified viral diseases (HIV, Herpes), V73.98 Special screening examination for unspecified chlamydial disease, V73.81 Special screening examination for Human papillomavirus (HPV)

A **32-year-old established female patient** presents seeking gonorrhea treatment. Her male partner of 1 year was treated 2 days prior after coming to the center for STI testing. She has been monogamous in the relationship. Using DMPA as contraception; condom use is described as "sporadic". Vitals taken and general appearance noted. GC /CT NAAT test ordered. Patient will be treated presumptively for gonorrhea. Ceftriaxone 250 mg IM injection given in right deltoid. Meds dispensed: Azithromycin 1GM po stat (dx V01.6). Patient is counseled face-to-face by the clinician for more than 50% of this 20-minute visit in regards to STI and safe sex practices. Time is documented. Patient will call for results. What codes do we need for this visit?

CPT Codes:

99213 for the problem-focused E/M for an established patient based on time, Outside lab will bill for ordered tests. Append a Modifier 25 to the E/M to show it is separate and distinct from the injection

Potential ICD-9 Codes:

V01.6 Contact with or exposure to venereal diseases, V73.98 Special screening examination for unspecified chlamydial disease

A **20-year-old single male** presents to your clinic requesting "Screen me for everything." He is an established patient. He has had multiple sexual partners, both male and female in the past 4 months including oral, anal (both insertive and receptive) and vaginal sex. He does not use condoms. No history of previous STI and is asymptomatic. Vitals taken, GU exam performed. Culture is done for gonorrhea (GC,) Chlamydia (CT), HPV, and herpes (HSV). Blood draw performed for syphilis (VDRL), HIV, hepatitis B and C. Patient instructed to return in 1 week for results. What codes do we need for this visit?

CPT Codes:

99213 Established patient problem focused E/M based on history, exam and medical decision-making, 36415 Venipuncture. Outside lab will bill for ordered tests

Potential ICD-9 Codes:

V69.2 High-risk sexual behavior, V74.5 Screening Venereal Disease, V73.89 Special screening for other specified viral diseases (HIV, Herpes), V73.98 Special screening examination for unspecified chlamydial disease, V73.81 Special screening examination for Human papillomavirus (HPV)

INSURANCE COMPANY INFORMATION

| Company | Contact Information |
|---------------------------------------|---------------------------------|
| AETNA | 1-800-US-AETNA (1-800-872-3862) |
| BLUE CROSS and BLUE SHIELD OF TEXAS | 1-800-451-0287 |
| CIGNA | 1-800-882-4462 (1-800-88CIGNA) |
| HUMANA | 1-800-4-HUMANA (1-800-448-6262) |
| UNITED HEALTHCARE | 1-866-633-2446 |
| NOVITAS (MEDICARE PART B-JH CONTRACT) | JH 1-855-252-8782 |

RESOURCES

- A guide to CIGNA's Preventive Health coverage for health care professionals. Cigna.com. July 2012. http://www.cigna.com/assets/docs/health-care-professionals/807467 d PreventiveHealthCovGuide v8 HR.pdf
- Aetna Clinical Policy Bulletin: HIV testing. http://www.aetna.com/cpb/medical/data/500 599/0542.html
- 3. American Medical Association, CPT. www.ama-assn.org/go/cpt
- 4. Centers for Disease Control and Prevention. Sexually Transmitted Diseases Treatment Guidelines, 2010. MMWR 2010; 59 (No. RR-12). http://www.cdc.gov/std/treatment/2010/STD-Treatment-2010-RR5912.pdf
- 5. Centers for Disease Control and Prevention. Revised Recommendations for HIV Testing of Adults, Adolescents, and Pregnant Women in Health-Care Settings. MMWR 2006; 55 (No. RR-14); 1-17. http://www.cdc.gov/mmwr/preview/mmwrhtml/rr5514a1.htm
- 6. Centers for Medicare and Medicaid (CMS). Evaluation and Management Services Guide.

 December 2010 / ICN: 006764. https://www.cms.gov/Outreach-and-Education/Medicare-Learning-Network-MLN/MLNProducts/downloads/eval_mgmt_serv_guide-ICN006764.pdf
- 7. HCPCS codes, Centers for Medicare and Medicaid services. <u>www.cms.gov/medhcpcsgeninfo</u>
- 8. Humana Preventive Services Guide, March 2011. http://zinserbenefitservice.com/wp-content/uploads/2011/04/Humana-Preventive-Services-Guide.pdf
- 9. ICD-9 Diagnosis Coding, ICD-9-CM Professional for Physicians- Volumes 1 & 2. OPTUM 2013.
- 10. ICD-10-CM, The Complete Official Draft Code Set. OPTUM 2013.
- Medicare Preventive Services MLN "Human Immunodeficiency Virus Screening" May 2011.
 http://www.cms.gov/Outreach-and-Education/Medicare-Learning-Network-MLN/MLNProducts/downloads/HIV brochure ICN905713.pdf
- 12. The Affordable Care Act: Preventive Services at 100%. BCBSTX, June 2012. http://www.tamus.edu/assets/files/benefits/pdf/programs/PreventiveHealthServicesFactSheet.pdf
- 13. United Healthcare Preventive Care Services: Coverage Determination Guideline CDG-A-036
 August 1, 2013. <a href="https://www.unitedhealthcareonline.com/ccmcontent/ProviderII/UHC/en-US/Assets/ProviderStaticFiles/ProviderStaticFilesPdf/Tools%20and%20Resources/Policies%20and%20Protocols/Medical%20Policies/Medical%20Policies/Preventive Care Services CD.pdf
- 14. US Preventive Task Force. Recommendations for STI Screening. http://www.uspreventiveservicestaskforce.org/uspstf08/methods/stinfections.htm