



Helpful HIV Medication Tables for Pharmacists

Northeast/Caribbean
AIDS Education
& Training Center (AETC)
www.nynjaetc.org

Summer 2016



Helpful HIV Medication Tables for Pharmacists


Dosing, Patient Counseling, and Drug Interactions

Developed by: John J. Faragon, PharmD, BCPS, AAHIVP,
Marshall J. Glesby, MD, PhD

This clinical support tool is sponsored by the Northeast/Caribbean AIDS Education Training Center (AETC). The Northeast/Caribbean AETC is funded by the Health Resources and Services Administration (HRSA) and is part of the National AIDS Education Training Center Program, a network of federally funded regional and national centers that conduct targeted multidisciplinary HIV/AIDS education and training programs for health care providers.

Disclaimer:

The data in this guide are intended for use by clinicians and other health care providers as guidance to minimize drug interactions and toxicities among patients being treated with HIV antiretroviral medications. These guidelines are for informational purposes only and cannot identify medical risks specific to an individual patient or recommend patient treatment. The absence of typographical errors is not guaranteed. These guidelines are not necessarily all-inclusive. Use of these guidelines indicates acknowledgement that neither Northeast/Caribbean AETC, nor the authors will be responsible for any loss or injury, sustained in connection with, or as a result of, the use of these guidelines. Users of this guide should consult other sources before prescribing medications or treatment. Data were compiled from Department of Health and Human Services Guidelines and product information for specific medications through Summer 2016.



Northeast/Caribbean AIDS Education and Training Center
Columbia University
College of Physicians and Surgeons
Department of Psychiatry
601 West 168th St, Suite 46
New York, NY 10032
(646) 774-6978
www.nynjaetc.org
nynjaetc@columbia.edu



Nucleoside/Nucleotide Reverse Transcriptase Inhibitors

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Abacavir (Ziagen®)

300mg twice daily or 600mg once daily

Food Effect – Take without regard to meals.

Adverse Effects – Patients should be warned about the abacavir hypersensitivity reaction (HSR) which is characterized by fever, rash, nausea, vomiting, malaise or fatigue, loss of appetite, sore throat, cough, shortness of breath.

Fatalities associated with the HSR have been reported, especially if patients are rechallenged.

HLA-B*5701 testing recommended prior to use. *

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Didanosine (Videx EC®)

**≥60kg – 400mg once daily; with tenofovir give 250mg once daily <60kg – 250mg once daily;
with tenofovir give 200mg once daily**

Food Effect – Take 1/2 hour before or 2 hours after a meal.

Adverse Effects – Peripheral neuropathy, pancreatitis and nausea. *

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Emtricitabine (Emtriva®)

200mg once daily

Food Effect – Take without regard to meals.

Adverse Effects – Minimal; Hyperpigmentation/skin discoloration has been reported. *

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Lamivudine (Epivir®)

150mg twice daily or 300mg once daily

Food Effect – Take without regard to meals.

Adverse Effects – Minimal; pancreatitis has been reported. *

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Stavudine (Zerit®)

≥60kg – 40mg twice daily <60kg – 30mg twice daily

Food Effect – Take without regard to meals.

Adverse Effects – Peripheral neuropathy, lipodystrophy, hyperlipidemia, pancreatitis. Rare, rapidly ascending neuromuscular weakness. *

* Lactic acidosis with hepatic steatosis is a rare, potentially life-threatening adverse event with the use of Nucleoside/Nucleotide Reverse Transcriptase Inhibitors.

Nucleoside/Nucleotide Reverse Transcriptase Inhibitors (Cont'd)

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Tenofovir (Viread®)

300mg once daily

Food Effect – Take without regard to meals.

Adverse Effects – Asthenia, headache, diarrhea, nausea, vomiting, flatulence, renal insufficiency. *

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Zidovudine (Retrovir®)

300mg twice daily or 200mg three times daily

Food Effect – Take without regard to meals.

Adverse Effects – Bone marrow suppression (macrocytic anemia, neutropenia), headache, insomnia, gastrointestinal intolerance, asthenia. *

* Lactic acidosis with hepatic steatosis is a rare, potentially life-threatening adverse event with the use of Nucleoside/Nucleotide Reverse Transcriptase Inhibitors.

Non-Nucleoside Reverse Transcriptase Inhibitors

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Delavirdine (Rescriptor®)

400mg three times daily

Food Effect – Take without regard to meals.

Adverse Effects – Rash, increased liver function tests, headache.

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Efavirenz (Sustiva®)

600mg once daily; preferably at bedtime

Food Effect – Take on an empty stomach.

Adverse Effects – Rash, central nervous system symptoms, lasting for approximately the first 2-4 weeks, including abnormal dreams, dizziness, somnolence and euphoria; increased liver function tests, false-positive cannabinoid test, teratogenic (Pregnancy Category D).

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Etravirine (Intence®)

200mg twice daily

Food Effect – Take after a meal. Fasting conditions reduce drug exposure by approximately 50%.

Adverse Effects – Rash (17%) and nausea. Stevens – Johnson Syndrome has been reported. Post marketing reports of fatalities due to toxic epidermal necrolysis, hypersensitivity reactions associated with liver failure have occurred.

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Nevirapine (Viramune®, Viramune XR®)

200mg once daily for 14 days, then 200mg twice daily or 400mg once daily if using XR

Food Effect – Take without regard to meals.

Adverse Effects – Rash, including Stevens-Johnson Syndrome; symptomatic hepatitis, including fatal hepatic necrosis reported. Higher frequency of hepatic events reported in treatment naïve females with CD4 >250 cells/mm³, and treatment naïve males with CD4 >400 cells/mm³.

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Rilpivirine (Edurant®)

25mg once daily

Food Effect – Take with a meal.

Adverse Effects – Depression, insomnia, headache, rash.

MEDICATION
STANDARD DOSING

Efavirenz/Tenofovir Disoproxil Fumarate/Emtricitabine (Atripla®)

Rilpivirine/Tenofovir Disoproxil Fumarate/Emtricitabine (Complera®)

Rilpivirine/Tenofovir Alafenamide/Emtricitabine (Odefsey®)

See Single Tablet Combination/Regimen in other section of guide

Protease Inhibitors

MEDICATION STANDARD DOSING PATIENT COUNSELING

Atazanavir (Reyataz®) and Atazanavir/cobicistat (Evotaz®)
400mg once daily or 300mg with ritonavir 100mg once daily or 300mg with cobicistat 150mg once daily

Food Effect – Take with food.

Adverse Effects – Indirect hyperbilirubinemia, nephrolithiasis, prolonged PR interval, (use with caution in patients with underlying conditions or concomitant medications that can cause PR prolongation); hyperglycemia, fat maldistribution, possible increased bleeding episodes in patients with hemophilia.

MEDICATION STANDARD DOSING PATIENT COUNSELING

Darunavir (Prezista®) and Darunavir/cobicistat (Prezcobix®)
**600mg with ritonavir 100mg twice daily or 800mg with ritonavir 100mg once daily
or 800mg with cobicistat 150mg once daily**

Food Effect – Take with food.

Adverse Effects – Skin rash (7%) including Stevens-Johnson Syndrome and erythema multiforme reported, caution in sulfa allergic patients, as darunavir contains a sulfonamide moiety; diarrhea, nausea, headache, hyperlipidemia, increased liver function tests, hepatotoxicity, hyperglycemia, fat maldistribution, possible increased bleeding episodes in patients with hemophilia.

MEDICATION STANDARD DOSING PATIENT COUNSELING

Fosamprenavir (Lexiva®)
1400mg twice daily or 1400mg with ritonavir 100 or 200mg once daily or 700mg with ritonavir 100mg twice daily

Food Effect – Take without regard to meals.

Adverse Effects – Skin rash (19%) including Stevens-Johnson Syndrome, caution in sulfa allergic patients, as fosamprenavir contains a sulfonamide moiety; diarrhea, nausea, vomiting, headache, hyperlipidemia, increased liver function tests, hyperglycemia, fat maldistribution, possible increased bleeding episodes in patients with hemophilia.

MEDICATION STANDARD DOSING PATIENT COUNSELING

Indinavir (Crixivan®)
800mg every 8 hours or 800mg with ritonavir 100mg every 12 hours

Food Effect – Requires 1.5 liters of fluid daily. Without ritonavir – Take 1 hour before or 2 hours after meals; may take with skim milk or low fat meal. With ritonavir – Take with or without food.

Adverse Effects – Nephrolithiasis, GI intolerance, nausea, indirect hyperbilirubinemia, hyperlipidemia, headache, asthenia, blurred vision, dizziness, rash, metallic taste, thrombocytopenia, alopecia, hemolytic anemia, hyperglycemia, fat maldistribution, possible increased bleeding episodes in patients with hemophilia.

Protease Inhibitors (Cont'd)

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Lopinavir/Ritonavir (Kaletra®)

Lopinavir 400mg/ritonavir 100mg twice daily or Lopinavir 800mg/ritonavir 200mg once daily

Food Effect – Take with food.

Adverse Effects – GI intolerance, nausea, vomiting, diarrhea, asthenia, hyperlipidemia (especially hypertriglyceridemia), increased liver function tests, hyperglycemia, fat maldistribution, possible increased bleeding episodes in patients with hemophilia.

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Nelfinavir (Viracept®)

1250mg twice daily or 750mg three times daily

Food Effect – Take with meal or snack. Levels increased 2-3 fold.

Adverse Effects – Diarrhea, hyperlipidemia, hyperglycemia, fat maldistribution, increased liver function tests, possible increased bleeding episodes in patients with hemophilia.

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Ritonavir (Norvir®)

100-200mg once or twice daily for protease inhibitor boosting depending on the protease inhibitor

Food Effect – Take with food to improve tolerability.

Adverse Effects – GI intolerance, nausea, vomiting, diarrhea, circumoral and extremity parasthesias, hyperlipidemia (especially hypertriglyceridemia), hepatitis, asthenia, taste perversion, hyperglycemia, fat maldistribution, possible increased bleeding episodes in patients with hemophilia.

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Saquinavir (Invirase®)

1000mg with ritonavir 100mg twice daily

Food Effect – Take within 2 hours of a meal when taken with ritonavir.

Adverse Effects – GI intolerance, nausea, diarrhea, headache, elevated liver function tests, hyperlipidemia, hyperglycemia, fat maldistribution, possible increased bleeding episodes in patients with hemophilia.

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Tipranavir (Aptivus®)

500mg with ritonavir 200mg twice daily

Food Effect – Take with food. High fat meals increase bioavailability.

Adverse Effects – Rash, caution in sulfa allergic patients, as tipranavir contains a sulfonamide moiety; hepatotoxicity including hepatic decompensation reported, especially in patients with underlying liver disease; hyperlipidemia, hyperglycemia, fat maldistribution, rare cases of fatal and non-fatal intracranial hemorrhages, possible increased bleeding episodes in patients with hemophilia.

Entry Inhibitors

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Enfuvirtide (Fuzeon®)

90mg SC twice daily

Adverse Effects – Local injection site reactions – pain, erythema, induration, nodules and cysts, pruritis, ecchymosis, bacterial pneumonia, hypersensitivity reaction (<1%) which includes rash, fever, nausea, vomiting, chills, rigors, hypotension, or increased liver function tests. Rechallenge not recommended.

**MEDICATION
STANDARD DOSING**

Maraviroc (Selzentry®)

**150mg twice daily when given with strong CYP3A inhibitors (with or without CYP3A inducers)
including PIs (except tipranavir/ritonavir)**

**300mg twice daily when given with NRTIs, enfuvirtide, tipranavir/ritonavir, nevirapine,
and other drugs that are not strong CYP3A inhibitors**

600mg twice daily when given with CYP3A inducers, including efavirenz, rifampin, etc. (without a CYP3A inhibitor)

PATIENT COUNSELING

Food Effect – Take with or without food.

Adverse Effects – Abdominal pain, cough, dizziness, musculoskeletal symptoms, pyrexia, rash, upper respiratory tract infections, hepatotoxicity, orthostatic hypotension.

Integrase Inhibitors

MEDICATION	Dolutegravir (Tivicay®)
STANDARD DOSING	50mg once daily or 50mg twice daily if integrase inhibitor experienced and resistance suspected or if combined with efavirenz, fosamprenavir/ritonavir, tipranavir/ritonavir or rifampin
PATIENT COUNSELING	Food Effect – Take with or without food. Adverse Effects – Insomnia, headache
MEDICATION	Elvitegravir, combined with Cobicistat, Tenofovir Disoproxil Fumarate and Emtricitabine (Stribild®)
STANDARD DOSING	See Single Tablet Combination/Regimen in other section of guide
MEDICATION	Elvitegravir, combined with Cobicistat, Tenofovir Alafenamide and Emtricitabine (Genvoya®)
STANDARD DOSING	See Single Tablet Combination/Regimen in other section of guide
MEDICATION	Dolutegravir, abacavir, lamivudine (Triumeq®)
STANDARD DOSING	See Single Tablet Combination/Regimen in other section of guide
MEDICATION	Raltegravir (Isentress®)
STANDARD DOSING	400mg twice daily
PATIENT COUNSELING	Food Effect – Take with or without food. Adverse Effects – Nausea, headache, diarrhea, pyrexia, rash, CPK elevation

Single Tablet Combinations and Regimens

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Efavirenz, Tenofovir, and Emtricitabine (Atripla®)

One tablet once daily, preferably at bedtime

Food Effect – Take on an empty stomach.

Adverse Effects – Rash, central nervous system symptoms, lasting for approximately the first 2-4 weeks, including abnormal dreams, dizziness, somnolence and euphoria; increased liver function tests, false-positive cannabinoid test, teratogenic (Pregnancy Category D), asthenia, headache, diarrhea, nausea, vomiting, flatulence, renal insufficiency, skin hyperpigmentation. *

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Elvitegravir, Cobicistat, Tenofovir Disoproxil Fumarate, Emtricitabine (Stribild®)

One tablet daily

Food Effect – Take with food.

Adverse Effects – Nausea, diarrhea, headache, elevated serum creatinine (artifact associated with cobicistat that is not a direct cause of renal impairment), renal impairment (associated with tenofovir)*

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Elvitegravir, Cobicistat, Tenofovir Alafenamide, Emtricitabine (Genvoya®)

One tablet daily

Food Effect – Take with food.

Adverse Effects – Nausea, diarrhea, headache, elevated serum creatinine (artifact associated with cobicistat that is not a direct cause of renal impairment). Tenofovir alafenamide has improved bone and renal safety profile compared to tenofovir disoproxil fumarate. *

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Dolutegravir, abacavir, lamivudine (Triumeq®)

One tablet daily

Food Effect – Take without regard to meals.

Adverse Effects – Patients should be warned about the abacavir hypersensitivity reaction (HSR), which is characterized by fever, rash, nausea, vomiting, malaise or fatigue, loss of appetite, sore throat, cough, shortness of breath. Fatalities associated with the HSR have been reported, especially if patients are rechallenged, rare pancreatitis. HLA-B*5701 testing recommended prior to use.*

* Lactic acidosis with hepatic steatosis is a rare, potentially life threatening adverse event with the use of Nucleoside/ Nucleotide Reverse Transcriptase Inhibitors.

Single Tablet Combinations and Regimens (Cont'd)

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Rilpivirine, Tenofovir Disoproxil Fumarate, and Emtricitabine (Complera®)

One tablet daily

Food Effect – Take with a meal.

Adverse Effects – Depression, insomnia, headache, diarrhea, nausea, vomiting, flatulence, renal insufficiency.*

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Zidovudine and Lamivudine (Combivir®)

One tablet twice daily

Food Effect – Take without regard to meals.

Adverse Effects – Bone marrow suppression (macrocytic anemia, neutropenia), headache, insomnia, gastrointestinal intolerance, asthenia, rare pancreatitis.*

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Abacavir and Lamivudine (Epzicom®)

One tablet once daily

Food Effect – Take without regard to meals.

Adverse Effects – Patients should be warned about the abacavir hypersensitivity reaction (HSR) which is characterized by fever, rash, nausea, vomiting, malaise or fatigue, loss of appetite, sore throat, cough, shortness of breath. Fatalities associated with the HSR have been reported, especially if patients are rechallenged, rare pancreatitis. HLA-B*5701 testing recommended prior to use.*

**MEDICATION
STANDARD DOSING
PATIENT COUNSELING**

Abacavir, Zidovudine, and Lamivudine (Trizivir®)

One tablet twice daily

Food Effect – Take without regard to meals.

Adverse Effects – Patients should be warned about the abacavir hypersensitivity reaction (HSR) which is characterized by fever, rash, nausea, vomiting, malaise or fatigue, loss of appetite, sore throat cough, shortness of breath. Fatalities associated with the HSR have been reported, especially if patients are rechallenged). HLA-B*5701 testing recommended prior to use. Bone marrow suppression (macrocytic anemia, neutropenia), headache, insomnia, gastrointestinal intolerance, asthenia, rare pancreatitis.

* Lactic acidosis with hepatic steatosis is a rare, potentially life threatening adverse event with the use of Nucleoside/Nucleotide Reverse Transcriptase Inhibitors.

Single Tablet Combinations and Regimens (Cont'd)

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Tenofovir Disoproxil Fumarate and Emtricitabine (Truvada®)

One tablet once daily

Food Effect – Take without regard to meals.

Adverse Effects – asthenia, headache, diarrhea, nausea, vomiting, flatulence, renal insufficiency, skin hyperpigmentation.*

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Tenofovir Alafenamide and Emtricitabine (Descovy®)

One tablet once daily

Food Effect – Take without regard to meals.

Adverse Effects – asthenia, headache, diarrhea, nausea, vomiting, flatulence, skin hyperpigmentation.

Tenofovir alafenamide has improved bone and renal safety profile compared to tenofovir disoproxil fumarate.*

MEDICATION
STANDARD DOSING
PATIENT COUNSELING

Rilpivirine/Tenofovir Alafenamide/Emtricitabine (Odefsey®)

One tablet once daily

Food Effect – Take with a meal.

Adverse Effects – Depression, insomnia, headache, diarrhea, nausea, vomiting, flatulence.

Tenofovir alafenamide has improved bone and renal safety profile compared to tenofovir disoproxil fumarate.*

* Lactic acidosis with hepatic steatosis is a rare, potentially life threatening adverse event with the use of Nucleoside/Nucleotide Reverse Transcriptase Inhibitors.

Select Medications to be Avoided with HIV Antiretroviral Therapy

MEDICATION OR CLASS	HIV MEDICATIONS TO BE AVOIDED
Afluzosin	Avoid with protease inhibitors and cobicistat
Antiarrhythmic	Avoid with fosamprenavir, saquinavir, and tipranavir. Use caution with other boosted protease inhibitors
Anticonvulsants	Avoid carbamazepine, oxcarbazepine, phenytoin and phenobarbital with dolutegravir, etravirine and rilpivirine, and with tenofovir alafenamide formulations.
Benzodiazepines – Midazolam and triazolam	Avoid with all protease inhibitors, cobicistat, delavirdine and efavirenz. Single doses of midazolam for sedation in controlled, monitored environment may be acceptable with intravenous midazolam. Oral midazolam should be avoided with protease inhibitors and cobicistat.
Ergot Alkaloids – Dihydroergotamine, ergotamine, ergonovine, methylegonovine	Avoid with all protease inhibitors, cobicistat, and all non-nucleoside reverse transcriptase inhibitors.
Fluticasone and budesonide	Avoid with all protease inhibitors and cobicistat, beclomethasone is a safe alternative.
Garlic supplements	Avoid with saquinavir
Irinotecan	Avoid with atazanavir and indinavir
Pimozide	Avoid with all protease inhibitors and cobicistat.
Proton pump inhibitors	Avoid with delavirdine, nelfinavir, and rilpivirine. With atazanavir, in treatment naïve patients, use only atazanavir 300mg with 100mg of ritonavir with a max dose equivalent to 20mg of omeprazole. Treatment experienced patients should not use proton pump inhibitors with unboosted or ritonavir boosted atazanavir. See atazanavir product information for additional dosing recommendations with proton pump inhibitors or H2 blockers.
Rifampin	Avoid with all protease inhibitors, cobicistat, delavirdine, etravirine and nevirapine, and tenofovir alafenamide formulations. Can be used with efavirenz; consider EFV dosage increase to 800mg daily. Can be used with raltegravir; increase raltegravir dosage to 800mg twice daily. Can be used with dolutegravir; increase dolutegravir dosage to 50mg twice daily
Salmeterol	Avoid with protease inhibitors
Sildenafil in pulmonary hypertension	Avoid with protease inhibitors and cobicistat. See DHHS Guidelines for more information.
St. John's Wort	Avoid with all Protease Inhibitors, Non nucleoside reverse transcriptase inhibitors, maraviroc, and cobicistat and with tenofovir alafenamide formulations.
Simvastatin and lovastatin	Avoid with all protease inhibitors, cobicistat and delavirdine

Components of an ARV Regimen Not Recommended in ANY Situation

REGIMEN/MEDICATION	RATIONALE
Atazanavir + Indinavir	Potential additive hyperbilirubinemia
Didanosine + Stavudine	High incidence of toxicities – peripheral neuropathy, pancreatitis, and hyperlactatemia. Reports of serious, even fatal, cases of lactic acidosis with hepatic steatosis with or without pancreatitis in pregnant women.
Dual Non Nucleoside Reverse Transcriptase Inhibitor combinations	Potential for higher incidence of adverse events; drug interactions complex and may lead to significant reductions in NNRTI drug levels
Efavirenz in first trimester or in women with significant childbearing potential	Teratogenic in humans and in nonhuman primates. Use only when no other antiretroviral options are available and potential benefits outweigh the risks.
Emtricitabine + lamivudine	Similar resistance profile, no potential benefit.
Etravirine + ritonavir boosted atazanavir, fosamprenavir, or tipranavir, OR any unboosted protease inhibitors	Etravirine may induce metabolism of protease inhibitors. Dosing with ATV, FPV, TPV not yet established.
Nevirapine initiation in treatment-naïve women with CD4 >250 cells/mm³ or in treatment-naïve men with CD4 >400 cells/mm³	Higher incidence of symptomatic (including serious and even fatal) hepatic events in these patient groups. Use only if the benefits clearly outweigh the risks.
Stavudine + zidovudine	Antagonistic effect on HIV-1
Unboosted darunavir, saquinavir, tipranavir	Poor oral bioavailability