

Pharmacy Information:

Clinic Information:

Clinical Team:

What are my HIV medication(s)?

What are my other medication(s)?

What is coinfection?

Coinfection occurs when a person with HIV is also infected with the hepatitis C virus (HCV).

HIV/HCV coinfection is a common scenario because of shared risk factors of both viruses.

HIV/HCV coinfection increases the risk of more severe liver disease, organ failure and other serious health issues.

Treatment of HCV in people with HIV should be a priority.

What kind of HCV do I have? (circled below)
There are 7 different HCV genotypes (types). A genotype is
the strain of the virus you have in your body.
You have genotype: 1a, 1b, 2, 3, 4, 5, 6
What stage of liver disease do I have?
Liver disease is "staged" based on the amount of scar tissue
or fibrosis in the liver: Mild Fibrosis (Stage 1) Cirrhosis (Stage 4)
□ Moderate Fibrosis (Stage 2) □ Decompensated Cirrhosis
Bridging Fibrosis (Stage 3) (Stage 5)
What is my HCV load after starting HCV meds?
0-4 Weeks
12 Weeks
24 Weeks
What was my last HIV viral load and CD4?
VL Date
CD4 Date
Which HCV medications am I on?
Daklinza® (daclatasvir) 1 tablet daily
🗅 90mg 🏾 🗅 60mg 🖉 🔄 30mg 🎽 🛛 🥮
Epclusa® (sofosbuvir/velpatasvir) 400mg/100mg
1 tablet daily
□ Harvoni [®] (ledipasvir/sofosbuvir) 90mg/400mg
1 tablet daily
Mavyret [®] (glecaprevir/pibrentasvir) 100mg/40mg
3 tablets daily with food
Copegus [®] , Rebetol [®] , Ribasphere [®] (ribavirin) 200mg
tablets twice daily with food
❑ Sovaldi [™] (sofosbuvir) 400mg
1 tablet daily
Uservi® (sofosbuvir/velpatasvir/voxilaprevir)
400mg/100mg/100mg 1 tablet daily with food
Zepatier [®] (elbasvir/grazoprevir) 50mg/100mg 1 tablet daily
How long is the HCV treatment?
■ 8 weeks ■ 12 weeks ■ 16 weeks ■weeks

This project is/was supported by the Health Resources and Services Administration (HRSA) of the U.S. Department of Health and Human Services (HHS) under grant number U10HA28686, AIDS Education and Training Center Program. This information or content and conclusions are those of the author and should not be construed as the official position or policy of, nor should any endorsements be inferred by HRSA, HHS or the U.S. Government. Rev Oct. 2019